VOL. 485 JOURNAL OF CHROMATOGRAPHY DECEMBER 27, 1989

CONTENTS

COMPUTER-ASSISTED METHOD DEVELOPMENT IN CHROMATOGRAPHY

Foreword
by J. L. Glajch and L. R. Snyder 1

Simplex optimization of high-performance liquid chromatographic separations (Review)
by J. C. Berridge (Sandwich, U.K.) 3

Computer-assisted optimization in high-performance liquid chromatographic method development
by S. N. Deming, J. M. Palasota, J. Lee and L. Sun (Houston, TX, U.S.A.) 15

Selection of mobile phase parameters and their optimization in reversed-phase liquid chromatography (Review)
by H. A. H. Billiet and L. de Galan (Delft, The Netherlands) 27

Method development in high-performance liquid chromatography using retention mapping and experimental design techniques (Review)
by J. L. Glajch (N. Billerica, MA, U.S.A.) and J. J. Kirkland (Wilmington, DE, U.S.A.) . 51

Drylab® computer simulation for high-performance liquid chromatographic method development. I. Isocratic elution (Review)
by L. R. Snyder, J. W. Dolan and D. C. Lommen (Lafayette, CA, U.S.A.) 65

Drylab® computer simulation for high-performance liquid chromatographic method development. II. Gradient elution (Review)
by J. W. Dolan, D. C. Lommen and L. R. Snyder (Lafayette, CA, U.S.A.) 91

Predictive calculation methods for optimization of gradient elution using binary and ternary solvent gradients
by P. Jandera (Pardubice, Czechoslovakia) 113

Computer-assisted retention prediction for high-performance liquid chromatography in the ion-exchange mode (Review)
by Y. Baba (Oita, Japan) 143

Multivariate calibration strategy for reversed-phase chromatographic systems based on the characterization of stationary–mobile phase combinations with markers
by A. K. Smilde, P. M. J. Coenegracht, C. H. P. Bruins and D. A. Doornbos (Groningen, The Netherlands) 169

Computer-aided optimization of high-performance liquid chromatography in the pharmaceutical industry (Review)
by E. P. Lankmayr and W. Wegscheider (Graz, Austria) and J. C. Gfeller, N. M. Djordjevic and B. Schreiber (Basle, Switzerland) 183

Comparison of optimization methods in reversed-phase high-performance liquid chromatography using mixture designs and multi-criteria decision making (Review)
by P. M. J. Coenegracht, A. K. Smilde, H. J. Metting and D. A. Doornbos (Groningen, The Netherlands) 195

Explanations and advice provided by an expert system for system optimization in high-performance liquid chromatography
by P. J. Schoenmakers (Eindhoven, The Netherlands) and N. Dunand (Cambridge, U.K.) 219

Expert system for the selection of high-performance liquid chromatographic methods for the analysis of drugs
by M. De Smet, G. Musch, A. Peeters, L. Buydens and D. L. Massart (Brussels, Belgium) 237

Expert system for the selection of initial high-performance liquid chromatographic conditions for the analysis of pharmaceuticals
by R. Hindriks, F. Maris and J. Vink (Oss, The Netherlands), A. Peeters, M. De Smet and D. L. Massart (Brussels, Belgium) and L. Buydens (Nijmegen, The Netherlands) 255

Expert system program for assistance in high-performance liquid chromatographic method development
by S. S. Williams, J. F. Karnicky, J.-L. Excoffier and S. R. Abbott (Walnut Creek, CA, U.S.A.) 267

Expert system for method validation in chromatography
by M. Mulholland, N. Dunand and A. Cleland (Cambridge, U.K.) and J. A. van Leeuwen and B. G. M. Vandeginste (Nijmegen, The Netherlands) 283

Knowledge-based expert system for troubleshooting high-performance liquid chromatographic assay methods
by K. Tsuji and K. M. Jenkins (Kalamazoo, MI, U.S.A.) 297

Uniform shell designs for optimization in reversed-phase liquid chromatography
by Y. Hu and D. L. Massart (Brussels, Belgium) 311

Retention prediction of analytes in reversed-phase high-performance liquid chromatography based on molecular structure. V. CRIPES (chromatographic retention index prediction expert system)
by R. M. Smith and C. M. Burr (Loughborough, U.K.) 325

Cathie: expert interpretation of chromatographic data
by R. Milne (Livingston Village, U.K.) 341

Prediction of retention of metabolites in high-performance liquid chromatography by an expert system approach
by K. Valkó (Budapest, Hungary), G. Szabó (Austin, TX, U.S.A.), J. Röhricht and K. Jemnitz (Budapest, Hungary) and F. Darvas (Austin,TX, U.S.A.) 349

Reversed-phase chromatographic method development for peptide separations using the computer simulation program ProDigest-LC
by C. T. Mant, T. W. L. Burke, N. E. Zhou, J. M.R. Parker and R. S. Hodges (Edmonton, Canada) 365

Rule-based approach for the determination of solute types in unknown sample mixtures as a first step of optimization parameter selection in reversed-phase ion-pair chromatography
by A. Bartha and G. Vigh (College Station, TX, U.S.A.) 383

Rationalization of the selection of the type of the organic modifier(s) for selectivity optimization in reversed-phase ion-pair chromatography
by A. Bartha and G. Vigh (College Station, TX, U.S.A.) and J. Ståhlberg (Södertälje, Sweden) 403

Predicting reversed-phase gradient elution separations by computer simulation. A comparison of two different programs
by J. Schmidt (Lexington, KY, U.S.A.) 421

Computer-assisted optimization with NEMROD software
by G. Mazerolles, D. Mathieu, R. Phan-Tan-Luu and A. M. Siouffi (Marseille, France) . 433

Multi-dimensional interpolation by the moving least-squares approach for modelling of chromatographic retention data
by M. Otto and U. Hörchner (Freiberg, G.D.R.) and W. Wegscheider (Graz, Austria) . . 453

Microcomputer-assisted liquid chromatographic separation system (MCASYST) for method development and data handling
by K. Jinno, M. Yamagami and M. Kuwajima (Toyohashi, Japan) 461

Objective functions in experimental and simulated chromatographic optimization. Comparative study and alternative proposals
by R. Cela (Santiago de Compostela, Spain) and C. G. Barroso and J. A. Pérez-Bustamante (Cádiz, Spain) 477

Optimization strategies for solutes exhibiting peak tailing. Comparison of two approaches
by S. Sekulic and P. R. Haddad (Kensington, Australia) 501

Computer-assisted selection of the optimum gradient programme in thin-layer chromatography
by W. Markowski (Lublin, Poland) 517

Prediction of retention times in ion-exchange chromatography
by T. Sasagawa, Y. Sakamoto, T. Hirose, T. Yoshida, Y.Kobayashi and Y. Sato (Kanagawa, Japan) and K. Koizumi (Nishinomiya, Japan) 533

Solvent modulation in liquid chromatography: optimization strategies
by J. H. Wahl and V. L. McGuffin (East Lansing, MI, U.S.A.) 541

Recent advances in fuzzy peak tracking in high-performance liquid chromatography
by E. P. Lankmayr, W. Wegscheider and J. Daniel-Ivad (Graz, Austria), I. Kolossvary and G. Csonka (Budapest, Hungary) and M. Otto (Freiberg, G.D.R.) 557

Peak tracking in high-performance liquid chromatography based on normalized band areas. A ribosomal protein sample as an example
by I. Molnar, R. Boysen and P. Jekow (Berlin, F.R.G.) 569

Development of a high-performance liquid chromatographic method for fluroxypyr herbicide and metabolites using computer simulation with Drylab G software
by R. G. Lehmann and J. R. Miller (Midland, MI, U.S.A.) 581

Computer-assisted optimization of a high-performance liquid chromatographic separation for chlorpromazine and thirteen metabolites
by J. S. Kiel, S. L. Morgan and R. K. Abramson (Columbia, SC, U.S.A.) 585

Practical approach for high-performance liquid chromatographic method development: assaying synthetic intermediates of a leukotriene inhibitor
by J. Fulper (Skokie, IL, U.S.A.) 597

Computer-assisted development of a high-performance liquid chromatographic method for fractionating selected nitro derivatives of polyaromatic hydrocarbons
by D. J. Thompson and W. D. Ellenson (Research Triangle Park, NC, U.S.A.) 607

Reversed-phase liquid chromatographic retention and selectivity surfaces. II. Deoxyribonucleosides
by E. Grushka (Jerusalem, Israel) and N.-I. Jang and P. R. Brown (Kingston, RI, U.S.A.) 617

Effects of different organic modifiers in optimization of reversed-phase high-performance liquid chromatographic gradient elution of a mixture of natural secoiridoid compounds
by F. Dondi (Ferrara, Italy), T. Gianferrara (Trieste, Italy), P. Reschiglian and M. C. Pietrogrande (Ferrara, Italy) and C. Ebert and P. Linda (Trieste, Italy) 631

Optimization of gradients in anion-exchange separations of oligonucleotides using computer-assisted retention prediction and a high-performance liquid chromatographic simulation system
by Y. Baba and M. K. Ito (Oita, Japan) 647

Separation of mixtures of *o*-phthalaldehyde-derivatized amino acids by reversed-phase gradient elution. Accuracy of computer simulation for predicting retention and band width
by J. D. Stuart and D. D. Lisi (Storrs, CT, U.S.A.) and L. R. Snyder (Orinda, CA, U.S.A.) 657

Author Index 673

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)